

On Riemann-Roch Formulas for Multiplicities

Eckhard Meinrenken ^{*†}

May 1994 [‡]

Abstract

A Theorem due to Guillemin and Sternberg [17] about geometric quantization of Hamiltonian actions of compact Lie groups G on compact Kähler manifolds says that the dimension of the G -invariant subspace is equal to the Riemann-Roch number of the symplectically reduced space. Combined with the shifting-trick, this gives explicit formulas for the multiplicities of the various irreducible components. One of the assumptions of the Theorem is that the reduction is regular, so that the reduced space is a smooth symplectic manifold. In this paper, we prove a generalization of this result to the case where the reduced space may have orbifold singularities. Our proof uses localization techniques from equivariant cohomology, and relies in particular on recent work of Jeffrey-Kirwan [20] and Guillemin [14]. Since there are no complex geometry arguments involved, the result also extends to non Kählerian settings.

1 Introduction

Let (M, ω) be a compact Kähler manifold, and let $\tau : L \rightarrow M$ be a holomorphic line bundle over M with Hermitian fiber metric h . (L, h) is said to satisfy the quantizing condition if $-2\pi i \omega$ is the curvature of the canonical Hermitian connection on L . Let $H^i(M, \mathcal{O}(L))$ be the i th cohomology group for the sheaf of germs of holomorphic sections. By the Riemann-Roch Formula of Hirzebruch and Atiyah-Singer[4], the Euler number

$$\text{Eul}(L) := \sum_i (-1)^i \dim H^i(M, \mathcal{O}(L)) \quad (1)$$

^{*}Mathematics Department, M.I.T., Cambridge MA 02139, U.S.A. Address after August 1, 1994: Fakultät für Physik, Hermann-Herder-Str. 3, D-79104 Freiburg, Germany.

[†]Supported by a grant from the German Academic Exchange Service.

[‡]Revised Version, October 1994

is equal to the characteristic number

$$\text{Eul}(L) = \int_M Td(M) Ch(L), \quad (2)$$

where $Td(M)$ is the Todd class and $Ch(L) = e^{[\omega]}$ the Chern character. Recall that if L is “sufficiently positive”, in particular if one replaces L by some sufficiently high tensor power, all the cohomology groups with $i > 0$ are zero by Kodaira’s Theorem [13], so in this case (2) gives a formula for the dimension of the space $H^0(M, \mathcal{O}(L)) = \Gamma_{hol}(M, L)$ of holomorphic sections.

Let G be a compact, connected Lie group that acts on M by Kähler diffeomorphisms $\Phi : G \times M \rightarrow M$, with an equivariant moment map $J : M \rightarrow \mathfrak{g}^*$. Suppose also that Φ lifts to Hermitian bundle automorphisms of $L \rightarrow M$, according to the rules of geometric quantization [17]. The corresponding virtual representation of G on $\sum (-1)^i H^i(M, \mathcal{O}(L))$ may then be regarded as the “quantization” of the classical action Φ . Its character χ is the element of the representation ring $R(G)$ defined by

$$\chi(g) := \sum_i (-1)^i \text{tr}(g|H^i(M, \mathcal{O}(L))). \quad (3)$$

From the Equivariant Riemann-Roch Formula of Atiyah-Segal-Singer [3, 4], one has an expression for $\chi(g)$ as the evaluation of certain characteristic classes on the fixed point set $M^g = \{x \in M | g.x = x\}$ (which is a Kähler submanifold of M): Let $Ch^g(L|M^g) = c_L(g) Ch(L|M^g)$, where $c_L(g) \in S^1$ is the (locally constant) action of g on $L|M^g$. Denote by N^g the normal bundle of M^g in M , by $F(N^g)$ its curvature, and let

$$D^g(N^g) = \det(I - (g^\sharp)^{-1} e^{-\frac{i}{2\pi} F(N^g)}) \quad (4)$$

where $g^\sharp$ is the automorphism of N^g defined by g . Then

$$\chi(g) = \int_{M^g} \frac{Td(M^g) Ch^g(L|M^g)}{D^g(N^g)}. \quad (5)$$

By a Theorem of Guillemin and Sternberg [17], there are also Riemann-Roch Formulas for the multiplicities of the irreducible components of the above representation, at least if certain regularity assumptions are satisfied. Let $T \subset G$ be a maximal torus, and $\mathfrak{g} = \mathfrak{t} \oplus [\mathfrak{t}, \mathfrak{g}]$ the corresponding decomposition of the Lie algebra. Choose a fundamental Weyl chamber $\mathfrak{t}_+^* \subset \mathfrak{t}^* \subset \mathfrak{g}^*$, let $\Lambda \subset \mathfrak{t}^*$ the integral lattice, and $\Lambda_+ = \Lambda \cap \mathfrak{t}_+^*$ the dominant weights. For a given lattice point $\mu \in \Lambda_+$, let V_μ denote the corresponding irreducible representation with highest weight μ , and define the multiplicity $N(\mu)$ by the alternating sum

$$N(\mu) := \sum_i (-1)^i \dim(V_\mu^* \otimes H^i(M, \mathcal{O}(L)))^G. \quad (6)$$

Suppose that $\mu \in \Lambda_+$ is a regular value of J , or equivalently that the action of the isotropy group G_μ on $J^{-1}(\mu)$ is locally free. If the action is in fact free, the reduced space $M_\mu = J^{-1}(\mu)/G_\mu$ is a smooth symplectic manifold, and it is well-known that it acquires a natural Kähler structure, together with a quantizing line bundle L_μ . The main result of [17] is that the multiplicity of μ in $\Gamma_{hol}(M, L)$ is equal to the dimension of the space $\Gamma_{hol}(M_\mu, L_\mu)$, so in particular $N(\mu)$ is given by the Euler number of L_μ if L is sufficiently positive:

Theorem 1.1 (V. Guillemin, S. Sternberg [17]) *If the action of G_μ on $J^{-1}(\mu)$ is free, and if L is sufficiently positive,*

$$N(\mu) = \int_{M_\mu} Td(M_\mu) Ch(L_\mu). \quad (7)$$

The “physical” interpretation of this Theorem is that reduction and quantization commute.

In practice, one is often dealing with situations where the action is only locally free. In this case, the reduced space is in general just an orbifold (or V-manifold) in the sense of Satake [22], which means (roughly) that it is locally the quotient of a manifold by a finite group. Moreover, the reduced line bundle is in general just an orbifold-bundle, that is, at some points the fiber of L_μ is not $\mathbb{C}$, but its quotient by a finite group. Guillemin and Sternberg conjectured that in this case, the right hand side of (7) has to be replaced by the expression from T. Kawasaki’s Riemann-Roch Formula for orbifolds [21]. It was proved by R. Sjamaar [23] that this assertion is true if L is sufficiently positive, and if L_μ is an honest line bundle. In fact, his approach also covers the truly singular case where μ is not even a regular value, by using Kirwan’s partial desingularization procedure to reduce it to the orbifold case. On the other hand, the condition that L_μ be a genuine line bundle is rather restrictive. It is the aim of the present paper to give a different proof of the Guillemin-Sternberg conjecture (for μ a regular value), without having to make this assumption.

The method we use is motivated by recent work of V. Guillemin [14], who used localization techniques from equivariant cohomology to establish the connection between the Multiplicity Formula (7) and a certain formula for counting lattice points in polytopes. This formula is known to be true in various interesting cases, and for these gives a new proof of (7) without using any complex geometry arguments. (In particular, it also works for *almost* Kähler polarizations.)

We will adopt this utilization of equivariant cohomology, but in a slightly different guise. The main idea is to consider the rescaled problem, where we replace

$$\omega \rightsquigarrow m\omega, L \rightsquigarrow L^m, J \rightsquigarrow mJ, \mu \rightsquigarrow m\mu \quad (8)$$

for $m \in \mathbb{N}$.

Our starting point will be the Equivariant Riemann-Roch Formula, but instead of (5) we will use it in a form due to Berline and Vergne [6], involving equivariant characteristic classes. By a stationary phase version of the Localization Formula of Jeffrey and Kirwan [20], we pass

from equivariant characteristic classes to (ordinary) characteristic classes on the reduced spaces. This leads to the desired Multiplicity Formula for $N^{(m)}(m\mu)$, up to an error term $O(m^{-\infty})$.

Since the multiplicities are integers, one easily finds that the error term is zero for large m . To investigate the general dependence of $N^{(m)}(m\mu)$ on m , we use a different expression for $N^{(m)}(m\mu)$ via the number of lattice points in certain polytopes. If $J(M)$ is contained in the set of regular elements of $\mathfrak{g}^*$, in particular in the abelian case, this analysis turns out to be sufficiently good to show that the above error term is zero for all m .

2 Statement of the result

In order to state the result, we have to give a closer description of the reduced space and its singular strata. Suppose that $\mu \in \Lambda_+$ is a regular value of J . Recall first the shifting trick to express M_μ as a reduced space at the zero level set: Let $O = G \cdot \mu$ be the coadjoint orbit through μ , equipped with its usual Kirillov Kähler structure, and let O^- denote O with the opposite Kähler structure. The action of G on O is Hamiltonian, with moment map Ψ the embedding into $\mathfrak{g}^*$. Then $\tilde{M} = M \times O^-$ is a Kähler manifold, and the diagonal action of G is Hamiltonian, with moment map $\tilde{J} = J - \Psi$. There are canonical identifications

$$M_\mu = J^{-1}(\mu)/G_\mu \cong J^{-1}(O)/G \cong \tilde{J}^{-1}(0)/G. \quad (9)$$

By Kostant's version of the Borel-Weil-Bott Theorem, one also has a natural quantizing bundle $\Xi \rightarrow O$, and the irreducible representation V_μ corresponding to μ gets realized as the space of holomorphic sections of Ξ . (The higher order cohomology groups $H^i(O, \mathcal{O}(\Xi))$ vanish.) The tensor product $\tilde{L} := L \otimes \Xi^*$ quantizes $\tilde{M}$, and there is an isomorphism

$$H^i(\tilde{M}, \mathcal{O}(\tilde{L})) \cong V_\mu^* \otimes H^i(M, \mathcal{O}(L)). \quad (10)$$

Hence $N(\mu) = \text{Eul}(\tilde{L})$, which is the quantum counterpart of the shifting-trick.

Using the shifting-trick, it is enough to consider the case $\mu = 0$. The reduced space M_0 inherits a natural Kähler structure from M (see [17]), and the reduced bundle $L_0 = (L|_{J^{-1}(0)})/G$ renders a quantizing orbifold-line bundle. Note however that L_0 need not be an honest line bundle, not even over the smooth part of M_0 . Sections of an orbifold bundle are defined as coming from invariant sections for the local orbifold charts, so all sections of L_0 have to vanish at points where the fiber is not $\mathbb{C}$.

Let us regard $P = J^{-1}(0)$ as an orbifold-principal bundle over $M_0 = J^{-1}(0)/G$. Following [21, 12], we introduce

$$\tilde{P} = \{(x, g) \mid x \in P, g \cdot x = x\} \subset P \times G, \quad (11)$$

and let $\Sigma = \tilde{P}/G$ be its quotient under the locally free action $h \cdot (x, g) = (h \cdot x, h g h^{-1})$. The projection of $\tilde{P}$ to the second factor descends to a locally constant mapping

$$\tau : \Sigma \rightarrow \text{Conj}(G) \quad (12)$$

to the set of conjugacy classes. For $g \in G$, let $(g) = \text{Ad}(G).g$ denote the corresponding conjugacy class, and Σ_g its preimage under τ . There is a natural identification $\Sigma_g = P^g/Z_g$, where $P^g \subset P$ is the fixed point manifold and Z_g the centralizer of g in G . Since the fixed point set $M^g \subset M$ is a Kähler submanifold, and the action of Z_g on M^g is Hamiltonian with the restriction of J serving as a moment map, this makes clear that Σ is a Kähler orbifold (with several components of different dimensions). Note that this Kähler structure does not depend on the choice of the representative for (g) . Observe also that $\Sigma_e \cong M_0$.

The collection of bundles $(L|P^g)/Z_g$ defines a quantizing orbifold line bundle $L_\Sigma \rightarrow \Sigma$. As above, let $c_L(g) \in S^1$ be the locally constant action of g on $L|P^g$, denote by $c_\Sigma : \Sigma \rightarrow S^1$ the function defined by the $c_L(g)$'s, and let $Ch^\Sigma(L_\Sigma)$ be the cohomology class defined by

$$Ch^\Sigma(L_\Sigma) = c_\Sigma e^{\omega_\Sigma} \quad (13)$$

where ω_Σ is the Kähler form on Σ .

Consider now the natural mapping $f : \Sigma \rightarrow M_0$, sending $G.(x, g) \rightarrow G.x$. In a local orbifold chart, the tangent space to Σ at $G.(x, g)$ is isomorphic to $T_x(M^g \cap J^{-1}(0))/T_x(Z_g.x)$, while the tangent space to M_0 at $G.x$ is $T_x(J^{-1}(0))/T_x(G.x)$. From this, it is easy to see that f is a Kähler immersion. Let $N_\Sigma \rightarrow \Sigma$ be the normal bundle of this immersion, and denote by $g^\sharp$ the automorphism of $N_\Sigma|_{\Sigma_g}$ induced by the action of g . Then the collection of differential forms

$$\det(I - (g^\sharp)^{-1} e^{-\frac{i}{2\pi} F(N_\Sigma)}), \quad (14)$$

where $F(N_\Sigma)$ is the curvature of N_Σ , defines a characteristic class $D^\Sigma(N_\Sigma)$.

Finally, for each connected component Σ_i of Σ , let d_i be the number of elements in a generic stabilizer for the G -action on the corresponding component $\tilde{P}_i$, and $d_\Sigma : \Sigma \rightarrow \mathbb{N}$ the function defined by the d_i 's.

For general values $\mu \in \Lambda_+$, let Σ_μ , L_μ etc. be defined by means of the shifting-trick. The main result of this paper is the following

Theorem 2.1 (Multiplicity Formula) *If $\mu \in \Lambda_+$ is a regular value of J , the multiplicities $N^{(m)}(m\mu)$ are for $m \gg 0$ given by the formula*

$$N^{(m)}(m\mu) = \int_{\Sigma_\mu} \frac{1}{d_{\Sigma_\mu}} \frac{Td(\Sigma_\mu) Ch^{\Sigma_\mu}(L_{\Sigma_\mu}^m)}{D^{\Sigma_\mu}(N_{\Sigma_\mu})}. \quad (15)$$

If the image of the moment map, $J(M)$, is contained in $\mathfrak{g}_{reg}^$ (the set of regular elements of $\mathfrak{g}^*$), one may set $m = 1$ in this formula:*

$$N(\mu) = \int_{\Sigma_\mu} \frac{1}{d_{\Sigma_\mu}} \frac{Td(\Sigma_\mu) Ch^{\Sigma_\mu}(L_{\Sigma_\mu})}{D^{\Sigma_\mu}(N_{\Sigma_\mu})}, \quad (16)$$

In particular, this is the case if G is abelian.

The following sections are aimed at proving this Theorem. We do not know whether the second part of the Theorem remains true without the condition $J(M) \subset \mathfrak{g}_{reg}^*$.

Remarks.

1. Comparing the right hand side of (16) to Kawasaki's Riemann-Roch Formula for orbifolds [21], the Theorem says that $N(\mu)$ is equal to the Euler number of the orbifold-bundle $L_\mu \rightarrow M_\mu$. In particular, $N(\mu)$ is zero if the fiber of L_μ over the smooth stratum of M_μ is a nontrivial quotient of $\mathbb{C}$.
2. Let $\Delta = J(M) \cap \mathfrak{t}_+^*$, which is a convex polytope by a result of Guillemin-Sternberg and Kirwan, and $\Delta^* \subset \Delta$ the set of regular values. By the Duistermaat-Heckman Theorem [9], the diffeotype of the reduced space M_μ (and of course also of Σ_μ) does not change as μ varies in a connected component of $int(\Delta^*)$, and the cohomology class of the symplectic form ω_μ varies linearly. In particular, the symplectic volume $Vol(M_\mu)$ is a polynomial on these connected components. If the action of G_μ on $J^{-1}(\mu)$ is free, so that $\Sigma_\mu = M_\mu$, the right hand side in (16) is equal to a polynomial as well, since all that varies is the Chern character $Ch(L_\mu) = e^{\omega_\mu}$. In the orbifold case, the behaviour is slightly more complicated: For $\mu \in \Lambda_+$ in any given connected component of $int(\Delta^*)$, and any connected component $\Sigma_{\mu,j}$ of Σ_μ ,

$$Ch^{\Sigma_\mu}(L_{\Sigma_\mu})|_{\Sigma_{\mu,j}} = \rho_\mu(g_j^{-1}) c_L(g_j) e^{\omega_{\Sigma_\mu}}, \quad (17)$$

where $g_j \in G_\mu$ represents $\tau(\Sigma_{\mu,j})$, and $\rho_\mu : G_\mu \rightarrow S^1$ is defined by $\rho_\mu(\exp(\xi)) = e^{2\pi i \langle \mu, \xi \rangle}$. Hence, the right hand side of (16) is of the form

$$N(\mu) = \sum_j \rho_\mu(g_j^{-1}) c_L(g_j) p_j(\mu), \quad (18)$$

where the p_j are polynomials of degree $\frac{1}{2} \dim(\Sigma_{\mu,j})$.

3. Since $Ch((L^m)_{m\mu}) = e^{m\omega_\mu}$, the right hand side of (15) is a polynomial in m if the G_μ -action on $J^{-1}(\mu)$ is free. In the orbifold-case, this is not true in general since

$$Ch^{\Sigma_\mu}(L_{\Sigma_\mu}^m)|_{\Sigma_{\mu,j}} = \rho_\mu(g_j^{-1})^m c_L(g_j)^m e^{m\omega_{\Sigma_\mu}}. \quad (19)$$

Definition 2.1 (Ehrhart [11]) *A function $f : \mathbb{N} \rightarrow \mathbb{C}$ is called an arithmetic polynomial, if for some $k \in \mathbb{N}$, all the functions*

$$q_j(m) = f(km - j), \quad j = 0, \dots, k-1 \quad (20)$$

are polynomials. k is called the period of f .

Equivalently, f is an arithmetic polynomial if and only if it can be written in the form

$$f(m) = \sum_{l=0}^{k-1} \exp(2\pi i \frac{l m}{k}) p_l(m), \quad (21)$$

where the p_l are polynomials. Taking k such that $g^k = e$ for all $(g) \in \tau(\Sigma_\mu)$, the right hand side of (15) clearly has this property.

4. Our proof of Theorem 2.1 does not really use the assumption that M is Kähler. Everything will be derived from the Equivariant Riemann-Roch Formula (3), which is of course valid in much more general situations. Suppose for instance that M is an arbitrary compact symplectic manifold, equipped with a Hamiltonian G -action, and that these data are quantizable. Then one can always choose a compatible, invariant almost Kähler structure, and replace the virtual space $\sum (-1)^i H^i(M, \mathcal{O}(L))$ with the index space of some G -invariant Dirac operator for the Clifford module $L \otimes \wedge(T^{(0,1)}M)^*$ (see [7, 14]). As an immediate consequence of the Berline-Vergne Formula for the character, Theorem 3.3 below, the multiplicities $N(\mu)$ defined in this way do not depend on the choice of the almost Kähler structure or of the quantizing line bundle L .

Example: Let $M = \mathbb{C}P(2)$, equipped with the Fubini-Study Kähler form ω_{FS} . Let $G = S^1$ act by

$$e^{i\phi} \cdot [z_0 : z_1 : z_2] = [e^{i\phi} z_0 : e^{-i\phi} z_1 : z_2].$$

This action is Hamiltonian, and has a moment map

$$J([z_0 : z_1 : z_2]) = \frac{|z_1|^2 - |z_0|^2}{|z_0|^2 + |z_1|^2 + |z_2|^2}.$$

The dual of the tautological line bundle serves as a quantizing line bundle L . We also consider the tensor powers L^m , which are quantizing line bundles for $(M, m\omega_{FS})$. By Kodaira's Theorem, $H^i(M, \mathcal{O}(L^m)) = 0$ for all $i > 0$, $m \in \mathbb{N}$. If we identify the spaces $\Gamma_{hol}(M, L^m)$ with the homogeneous polynomials of degree m on $\mathbb{C}^3$, the representation of S^1 is induced by the action $e^{i\phi} \cdot (z_0, z_1, z_2) = (e^{i\phi} z_0, e^{-i\phi} z_1, z_2)$ on $\mathbb{C}^3$. The isotypical subspace of $\Gamma_{hol}(M, L^m)$ corresponding to the weight $l \in \mathbb{Z}$ is, for $l \geq 0$, spanned by

$$z_0^l z_2^{m-l}, z_0^{l+1} z_1 z_2^{m-l-2}, \dots, z_0^{l+r} z_1^r z_2^{m-l-2r}$$

with $r = \lfloor \frac{m-l}{2} \rfloor$. For $l \leq 0$, the roles of z_0 and z_1 are reversed. Thus

$$N^{(m)}(l) = \begin{cases} 1 + \lfloor \frac{m-|l|}{2} \rfloor & \text{if } |l| \leq m \\ 0 & \text{otherwise} \end{cases},$$

for all $l \in \mathbb{Z}$, $m \in \mathbb{N}$. On the other hand, the image of the moment map $J^{(m)} = mJ$ is the interval $-m \leq \mu \leq m$, with critical values at $-m, 0, m$. If $0 < |l| < m$, the level set $(mJ)^{-1}(l)$ consists of two orbit type strata: On the set where $z_2 \neq 0$, the action is free, and on the set where $z_2 = 0$, the stabilizer is $\mathbb{Z}_2$. Writing $S = \{e, g\}$, the reduced space $M_\mu^e = M_\mu$ is an orbifold with a $\mathbb{Z}_2$ singularity (the “teardrop-orbifold”), and M_μ^g is the singular point.

Since $c_{L^m}(g) = (-1)^m$ and $\rho_l(g^{-1}) = (-1)^l$, we expect the multiplicities to grow like $N^{(m)}(l) = p_e(m, l) + (-1)^{(m-l)}p_g(m, l)$ where p_e is a first order polynomial and p_g a constant. Comparison with the above formula shows that this is indeed the case, with $p_e(m, l) = 3/4 + (m - |l|)/2$ and $p_g(m, l) = 1/4$. Note that for $m - |l|$ even, the fiber of L_l^m at the singular point is $\mathbb{C}$, whereas for $m - |l|$ odd it is $\mathbb{C}/\mathbb{Z}_2$. This means that for $m - |l|$ odd, all holomorphic sections of L_l^m have to vanish at the singular point. Again, this fits with the above explicit formulas.

3 Some equivariant cohomology

We start by reviewing Cartan’s model for equivariant cohomology, following Berline and Vergne [6]. Let M be a compact manifold, G a compact Lie group, and $\Phi : G \times M \rightarrow M$ a smooth action. Denote by $\mathcal{A}_G(M)$ the space of G -invariant polynomial mappings $\sigma : \mathfrak{g} \rightarrow \mathcal{A}(M)$, that is, $\sigma(\xi)$ depends polynomially on ξ and satisfies the equivariance property

$$\sigma(\text{Ad}_g(\xi)) = \Phi_{g^{-1}}^*(\sigma(\xi)). \quad (22)$$

The elements of $\mathcal{A}_G(M)$ are called *equivariant differential forms*, and the space $\mathcal{A}_G(M)$ is preserved under the *equivariant differential*

$$d_{\mathfrak{g}} : \mathcal{A}_G(M) \rightarrow \mathcal{A}_G(M), \quad (d_{\mathfrak{g}}\sigma)(\xi) = d(\sigma(\xi)) + 2\pi i(\iota(\xi_M)\sigma(\xi)). \quad (23)$$

Here, ξ_M denotes the fundamental vector field, i.e. the generating vector field of the flow $(t, p) \mapsto \exp(-t\xi).p$.

Equivariance together with Cartan’s identity for the Lie derivative, $\mathcal{L}_Y = \iota_Y \circ d + d \circ \iota_Y$, implies $d_{\mathfrak{g}}^2 = 0$. The cohomology $H_G(M)$ of the complex $(\mathcal{A}_G(M), d_{\mathfrak{g}})$ is called the *equivariant cohomology*. One can show that if the action of G on M is locally free, the pullback mapping $\mathcal{A}(M/G) \rightarrow \mathcal{A}(M)_{hor}^G \hookrightarrow \mathcal{A}_G(M)$ gives rise to an isomorphism $H(M/G) \rightarrow H_G(M)$. After choosing a principal connection on $M \rightarrow M/G$, the inverse is induced on the level of forms by the mapping

$$\mathcal{A}_G(M) \rightarrow \mathcal{A}(M)^G \rightarrow \mathcal{A}(M)_{hor}^G \cong \mathcal{A}(M/G) \quad (24)$$

given by substituting $\frac{i}{2\pi}$ times the curvature in the $\mathfrak{g}$ -slot, followed by projection onto the horizontal part (for the proof, see [10]).

For what follows, it will be necessary to relax the polynomial dependence on ξ to analytic dependence, possibly defined only on some neighborhood of $0 \in \mathfrak{g}$. We will denote the corresponding space of equivariant forms by $\mathcal{A}_G^\omega(M)$, and the cohomology by $H_G^\omega(M)$.

Suppose now that $\mathcal{V} \rightarrow M$ is a G -equivariant Hermitian vector bundle over M , with fiber dimension N . Let $\mathcal{A}(M, \mathcal{V})$ be the bundle-valued differential forms, and $\mathcal{A}_G(M, \mathcal{V})$ their equivariant counterpart. For each G -invariant Hermitian connection $\nabla : \mathcal{A}(M, \mathcal{V}) \rightarrow \mathcal{A}(M, \mathcal{V})$, the moment map $\mu \in \mathcal{A}_G(M, \text{End}(\mathcal{V}))$ of Berline and Vergne is defined by

$$\mu(\xi) \cdot \sigma := \xi \cdot \sigma - \nabla_{\xi_M} \sigma, \quad (25)$$

where $\sigma \rightarrow \xi \cdot \sigma$ denotes the representation of $\mathfrak{g}$ on the space of sections. Geometrically, $\mu(\xi)$ is the vertical part (with respect to the connection) of the fundamental vector field $\xi_{\mathcal{V}}$ on $\mathcal{V}$. Let $F(\mathcal{V}) \in \mathcal{A}^2(M, \text{End}(\mathcal{V}))$ denote the curvature of ∇ . The *equivariant* curvature is then defined by

$$F_{\mathfrak{g}}(\mathcal{V}, \xi) = F(\mathcal{V}) + 2\pi i \mu(\xi), \quad (26)$$

and it satisfies the Bianchi identity with respect to the equivariant covariant derivative $\nabla_{\mathfrak{g}} = \nabla + 2\pi i \iota(\xi_M)$. Suppose now that $A \rightarrow f(A)$ is the germ of a $U(N)$ -invariant analytic function on $\mathfrak{u}(N)$. Then $f(F_{\mathfrak{g}}) \in \mathcal{A}_G(M)$ is $d_{\mathfrak{g}}$ -closed, and one can show that choosing a different connection changes $f(F_{\mathfrak{g}})$ by a $d_{\mathfrak{g}}$ -exact form. The corresponding cohomology classes are called the *equivariant characteristic classes* of $\mathcal{V} \rightarrow M$. If the action on M is locally free, one can choose ∇ in such a way that $\mu = 0$, which shows that the mapping $H_G^\omega(M) \rightarrow H(M/G)$ sends the equivariant characteristic classes of $\mathcal{V}$ to the usual characteristic classes of the orbifold-bundle $\mathcal{V}/G$.

The following characteristic classes will play a role in the sequel:

- (a) The equivariant Chern character, defined by

$$Ch_{\mathfrak{g}}(\mathcal{V}, \xi) = \text{tr}(e^{\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi)}). \quad (27)$$

In the above geometric quantization setting, $\mathcal{V} = L$ is a line bundle, and for the equivariant curvature one has $\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi) = \omega + 2\pi i \langle J, \xi \rangle$, thus

$$Ch_{\mathfrak{g}}(L, \xi) = e^{\omega + 2\pi i \langle J, \xi \rangle}. \quad (28)$$

More generally, if $g \in G$ acts trivially on the base M , one defines

$$Ch_{\mathfrak{g}}^g(\mathcal{V}, \xi) = \text{tr}(g^{\mathcal{V}} e^{\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi)}) \quad (29)$$

where $g^{\mathcal{V}} \in \Gamma(M, \text{End}(\mathcal{V}))$ is the induced action of g . In the line bundle case, this is simply $c_L(g) Ch_{\mathfrak{g}}(L, \xi)$ where $c_L(g) \in S^1$ is the action of g on the fibers.

(b) The equivariant Todd class,

$$Td_{\mathfrak{g}}(\mathcal{V}, \xi) = \det \left(\frac{\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi)}{1 - e^{-\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi)}} \right). \quad (30)$$

The Todd class of a complex manifold is defined as the Todd class of its tangent bundle.

(c) The equivariant Euler class

$$\chi_{\mathfrak{g}}(\mathcal{V}, \xi) = \det \left(\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi) \right). \quad (31)$$

(d) The class

$$D_{\mathfrak{g}}^g(\mathcal{V}, \xi) = \det(I - (g^{-1})^{\mathcal{V}} e^{-\frac{i}{2\pi} F_{\mathfrak{g}}(\mathcal{V}, \xi)}), \quad (32)$$

for $g \in G$ acting trivially on M .

All of this also makes sense for symplectic vector bundles, since the choice of a compatible complex structure reduces the structure group to $U(n)$, and any two such choices are homotopic.

The equivariant Euler class occurs in the Localization Formula of Atiyah-Bott and Berline-Vergne.

Theorem 3.1 (Atiyah-Bott [2], Berline-Vergne [5]) *Suppose M is an orientable G -manifold, and $\sigma \in \mathcal{A}_G^{\omega}(M)$ is $d_{\mathfrak{g}}$ -closed. Assume that $\xi \in \mathfrak{g}$ is in the domain of definition of $\sigma(\xi)$, and let F be the set of zeroes of ξ_M . The connected components of F are then smooth submanifolds of even codimension, and the normal bundle N_F admits a Hermitian structure which is invariant under the flow of ξ_M . Choose orientations on F and M which are compatible with the corresponding orientation of N_F . Then*

$$\int_M \sigma(\xi) = \int_F \frac{\iota_F^* \sigma(\xi)}{\chi_{\mathfrak{g}}(N_F, \xi)}, \quad (33)$$

where $\iota_F : F \rightarrow M$ denotes the embedding.

We will need this result only in the symplectic or complex case, where the above orientations are given in a natural way.

Jeffrey and Kirwan [20] have proved a different sort of Localization Formula for the case of *Hamiltonian* G -spaces. We will need a stationary phase version of their result. Let (M, ω) be a Hamiltonian G -space, with moment map $J : M \rightarrow \mathfrak{g}^*$, and suppose 0 is a regular value of J . Let $\sigma \in \mathcal{A}_G^{\omega}(M)$ be $d_{\mathfrak{g}}$ -closed, and let $\Delta \in C_0^{\infty}(\mathfrak{g})$ be a cutoff-function, with $\sigma(\xi)$ defined for $\xi \in \text{supp}(\Delta)$, and $\Delta = 1$ in a neighborhood of 0. Consider the integral

$$\int_{\mathfrak{g}} \int_M \Delta(\xi) \sigma(\xi) e^{m(\omega + 2\pi i \langle J, \xi \rangle)} d\xi, \quad (34)$$

where $d\xi$ is the measure on $\mathfrak{g}$ corresponding to the normalized measure on G . Notice that the “equivariant symplectic form” $\omega + 2\pi i \langle J, \xi \rangle$ is $d_{\mathfrak{g}}$ -closed. Since $e^{m\omega}$ is simply a polynomial in m , the leading behaviour of this integral for $m \rightarrow \infty$ is determined by the stationary points of the phase function $e^{2\pi i m \langle J, \xi \rangle}$. Stationarity in $\mathfrak{g}$ -direction gives the condition $J = 0$, and stationarity in M -direction the condition $d\langle J, \xi \rangle = 0$, or $\xi = 0$ since the action on $J^{-1}(0)$ is locally free. Let $M_0 = J^{-1}(0)/G$ be the reduced space, $\pi : J^{-1}(0) \rightarrow M_0$ the projection and $\iota : J^{-1}(0) \rightarrow M$ the embedding. Consider the mapping

$$\kappa : H_G^\omega(M) \rightarrow H(M_0), \quad (35)$$

given by composing pullback to $J^{-1}(0)$ with the mapping $H_G^\omega(J^{-1}(0)) \rightarrow H(M_0)$. On the level of forms $\sigma \in \mathcal{A}_G^\omega(M)$, the form $\pi^* \kappa(\sigma)$ is by definition equal to the horizontal part of $\iota^* \sigma(\frac{i}{2\pi} F^\theta)$, where $F^\theta \in \mathcal{A}^2(J^{-1}(0), \mathfrak{g})_{hor}^G$ is the curvature of some connection $\theta \in \mathcal{A}^1(J^{-1}(0), \mathfrak{g})^G$.

Theorem 3.2 *For $m \rightarrow \infty$,*

$$\int_{\mathfrak{g}} \int_M \Delta(\xi) \sigma(\xi) e^{m(\omega + 2\pi i \langle J, \xi \rangle)} d\xi = \frac{1}{d} \int_{M_0} \kappa(\sigma) e^{m\omega_0} + O(m^{-\infty}), \quad (36)$$

where d is the number of elements in the generic stabilizer for the G -action on $J^{-1}(0)$.

Proof.

This is a variation of Theorem 4.1 in Jeffrey-Kirwan [20], which deals with polynomial equivariant cohomology classes, and where one has a Gaussian convergence factor instead of the cutoff.

Following [20], we will perform the integral in a local model for M near $J^{-1}(0)$. By the Coisotropic Embedding Theorem of Gotay, a neighborhood of $J^{-1}(0)$ in M is equivariantly symplectomorphic to a neighborhood of the zero section of the trivial bundle $J^{-1}(0) \times \mathfrak{g}^*$, with symplectic form $\pi^* \omega_0 + d\langle \alpha, \theta \rangle$, where α is the coordinate function on $\mathfrak{g}^*$. In this model, G acts by $g.(x, \alpha) = (g.x, \text{Ad}_{g^{-1}}^* \alpha)$, and the moment map is simply $J(x, \alpha) = \alpha$. Using the model and another cutoff-function $\Delta'(\alpha)$ on $\mathfrak{g}^*$, equal to 1 near the origin and with sufficiently small support, the same computation as in [20] shows that the integral is equal to

$$m^{\dim G} \int_{J^{-1}(0)} \int_{\mathfrak{g}^*} \int_{\mathfrak{g}} \Delta(\xi) \Delta'(\alpha) (\iota^* \sigma)(\xi) e^{m(\pi^* \omega_0 + \langle \alpha, F^\theta + 2\pi i \xi \rangle)} d\xi d\alpha dg + O(m^{-\infty}).$$

Here, dg denotes the (vertical) volume form on the fibers of $J^{-1}(0) \rightarrow M_0$, corresponding to the canonical identification $T_x(\text{fiber}) \cong \mathfrak{g}$ by means of the G -action.

Now apply the Stationary Phase Theorem (see e.g. [19], Theorem 7.7.5) to the α, ξ -integral, the relevant phase function being $e^{2\pi i m \langle \alpha, \xi \rangle}$. Since $e^{m \langle \alpha, F^\theta \rangle}$ is simply a polynomial in α , the

stationary phase expansion terminates after finitely many terms, and the result is

$$\begin{aligned} \int_{J^{-1}(0)} e^{m\pi^*\omega_0} \sum_{r=0}^{\infty} \frac{1}{r!} \left(\frac{i}{2\pi m} \right)^r \left(\sum_j \frac{\partial}{\partial \xi_j} \frac{\partial}{\partial \alpha^j} \right)^r \bigg|_{\substack{\xi=0 \\ \alpha=0}} \iota^* \sigma(\xi) e^{m\langle \alpha, F^\theta \rangle} dg + O(m^{-\infty}) \\ = \int_{J^{-1}(0)} \pi^* (e^{m\omega_0}) \iota^* \sigma \left(\frac{i}{2\pi} F^\theta \right) dg + O(m^{-\infty}). \end{aligned}$$

Since $\iota^* \sigma(\frac{i}{2\pi} F^\theta)$ gets wedged with dg , only its horizontal part, which by definition of κ is $\pi^* \kappa(\sigma)$, contributes to the integral. The result (36) now follows by integration over the fiber; the factor $1/d$ appears since this is the volume of a generic fiber. $\square$

We now turn to the Equivariant Hirzebruch-Riemann-Roch Theorem, in the form due to Berline and Vergne [6]. Let M be a compact complex manifold, equipped with a holomorphic action of G , and let $L \rightarrow M$ be a G -equivariant holomorphic line bundle. Define the character $\chi \in R(G)$ as in (3).

Theorem 3.3 *For ξ sufficiently close to zero,*

$$\chi(e^\xi) = \int_M Td_{\mathfrak{g}}(M, \xi) Ch_{\mathfrak{g}}(L, \xi). \quad (37)$$

More generally, if $g \in G$, one has for all sufficiently small $\xi \in \mathfrak{k}$, the Lie algebra of the centralizer Z_g of g :

$$\chi(g e^\xi) = \int_{M^g} \frac{Td_{\mathfrak{k}}(M^g, \xi) Ch_{\mathfrak{k}}^g(L|_{M^g}, \xi)}{D_{\mathfrak{k}}^g(N^g, \xi)}, \quad (38)$$

where M^g is the fixed point set and $N^g \rightarrow M^g$ its normal bundle.

To be precise, Berline and Vergne have shown how to rewrite the Equivariant Atiyah-Segal-Singer Index Theorem for Dirac operators in this style, with the equivariant $\hat{A}$ -genus appearing on the right hand side. This formula, however, implies the above Theorem in the same way as the usual Atiyah-Singer Index Theorem leads to the Hirzebruch-Riemann-Roch Formula; see [7], p. 152 for the calculations.

4 The stationary phase approximation

In this section, we will prove the first part of Theorem 2.1.

By the shifting trick, it is sufficient to consider the case $\mu = 0$. The idea is to substitute the expressions from the Equivariant Hirzebruch-Riemann-Roch Theorem 3.3 for $\chi^{(m)}$ in

$$N^{(m)}(0) = \int_G \chi^{(m)}(h) dh, \quad (39)$$

and apply the Localization Formula, Theorem 3.2. For this, we need to know what happens to the equivariant Todd class of M under the mapping (35):

Lemma 4.1 *Let*

$$j_{\mathfrak{g}}(\xi) = \det \left(\frac{1 - e^{-\text{ad}(\xi)}}{\text{ad}(\xi)} \right) \quad (40)$$

be the Jacobian of the exponential mapping $\exp : \mathfrak{g} \rightarrow G$. Then

$$\kappa(Td_{\mathfrak{g}}(M)j_{\mathfrak{g}}) = Td(M_0). \quad (41)$$

Proof. Identify the vertical subbundle of $TJ^{-1}(0)$ with the trivial bundle $\mathfrak{g}$, and the symplectic bundle $\mathfrak{g} \oplus I\mathfrak{g}$ (where I is the complex structure of M) with $\mathfrak{g}_{\mathbb{C}}$. Then

$$\iota^*(TM) = \pi^*(TM_0) \oplus \mathfrak{g}_{\mathbb{C}}.$$

Since the equivariant Todd class of $\mathfrak{g}_{\mathbb{C}}$ is just $j_{\mathfrak{g}}^{-1}$, this shows $\kappa(Td_{\mathfrak{g}}(M)) = Td(M_0)\kappa(j_{\mathfrak{g}}^{-1})$, q.e.d. $\square$

We will now consider the contribution to (39) coming from a small Ad -invariant neighborhood of a given orbit $(g) = Ad(G).g$. Let $\sigma \in C^\infty(G)$ be an Ad -invariant cutoff-function, supported in a sufficiently small neighborhood of (g) and equal to 1 near (g) . Consider the integral

$$I_g(m) = \int_G \sigma(h) \chi^{(m)}(h) dh. \quad (42)$$

Since (38) only holds for $\xi \in \mathfrak{k}$, we want to replace this integral by an integral over Z_g . (Of course, this step is void in the abelian case.)

Let $\mathfrak{r} \subset \mathfrak{g}$ be the orthogonal of $\mathfrak{k}$ with respect to some invariant inner product (or, more intrinsically, the annihilator of $(\mathfrak{g}^*)^g \cong \mathfrak{k}^*$). For $k \in Z_g$, let $k^{\mathfrak{r}}$ denote the action of k on $\mathfrak{r}$.

Lemma 4.2 *Let $f \in C^\infty(G)$ be $Ad(G)$ -invariant, with support in a small neighborhood of $Ad(G).g$. Then, for a suitable $Ad(Z_g)$ -invariant cutoff-function $\tilde{\sigma} \in C_0^\infty(Z_g)$, supported near $e \in Z_g$ and identically 1 near e ,*

$$\int_G f(h) dh = \int_{Z_g} f(gk) \det(I - (g^{-1}k^{-1})^{\mathfrak{r}}) \tilde{\sigma}(k) dk. \quad (43)$$

Proof. From the Slice Theorem for actions of compact Lie groups, it follows that an invariant neighborhood of the orbit $Ad(G).g$ is equivariantly diffeomorphic to a neighborhood of the zero section of $G \times_{Z_g} \mathfrak{k}$, where Z_g acts on $\mathfrak{k}$ by the adjoint action. Using the exponential map for Z_g , it follows that the mapping

$$\phi : G \times_{Z_g} Z_g \rightarrow G, (h, k) \mapsto Ad_h(gk), \quad (44)$$

with Z_g acting on itself by Ad , is an equivariant diffeomorphism from a neighborhood of the unit section to a neighborhood of the orbit. Let $d\nu$ be the canonical measure on the group bundle $G \times_{Z_g} Z_g =: W$, constructed from the normalized invariant measures on Z_g and G/Z_g . We have to compute the tangent mapping to ϕ , but by equivariance it is sufficient to do this at points (e, k) in the fiber over $e Z_g$, which is canonically isomorphic to Z_g . If we identify $T_{(e,k)}W \cong \mathfrak{r} \oplus \mathfrak{k}$, and $TG \cong G \times \mathfrak{g}$ using left trivialization, the tangent mapping is given by

$$T_{(e,k)}\phi(\xi, \eta) = ((1 - \text{Ad}_{(gk)^{-1}})(\xi), \eta).$$

This shows that the measure transforms according to

$$\phi^* dg = \det(I - (g^{-1}k^{-1})^{\mathfrak{r}}) d\nu. \quad (45)$$

We can now perform the integral by first pulling f back to $G \times_{Z_g} Z_g$, multiplying with a suitable cutoff function $\tilde{\sigma}$, integrating over the fibers of $W \rightarrow G/Z_g$, and then integrate over the base $\text{Ad}(G).g \cong G/Z_g$. $\square$

Using the Lemma, we find that

$$I_g(m) = \int_{Z_g} \tilde{\sigma}(k) \chi^{(m)}(gk) \det(I - (g^{-1}k^{-1})^{\mathfrak{r}}) dk. \quad (46)$$

Replacing this with an integral over the Lie algebra $\mathfrak{k}$, and using (38) gives

$$I_g(m) = \int_{\mathfrak{k}} \int_{M^g} \Delta(\xi) \frac{Td_{\mathfrak{k}}(M^g, \xi) Ch_{\mathfrak{k}}^g(L^m | M^g, \xi)}{D_{\mathfrak{k}}^g(N^g, \xi)} \det(I - (g^{-1}e^{-\xi})^{\mathfrak{r}}) j_{\mathfrak{k}}(\xi) d\xi, \quad (47)$$

with $\Delta(\xi) = \tilde{\sigma}(e^{\xi})$. Let $\kappa_g : H_{Z_g}(M^g) \rightarrow H(\Sigma_g)$ be the mapping defined by (35), with M replaced by M^g and G by Z_g . By Lemma 4.1,

$$\kappa_g(Ch_{\mathfrak{k}}^g(L^m | M^g)) \kappa_g(j_{\mathfrak{k}}) = Ch^{\Sigma}(L_{\Sigma}^m) | \Sigma_g. \quad (48)$$

For $x \in P^g = M^g \cap J^{-1}(0)$, let $\mathfrak{r}_M(x) := \{\xi_M(x) | \xi \in \mathfrak{r}\} \cong \mathfrak{r}$. Then

$$N^g(x) = N_{\Sigma}(y) \oplus \mathfrak{r}_M(x) \oplus I\mathfrak{r}_M(x) = N_{\Sigma}(y) \oplus \mathfrak{r}_{\mathbb{C}}, \quad (49)$$

where $y = G.(x, g)$. But $D_{\mathfrak{k}}^g(\mathfrak{r}_{\mathbb{C}}, \xi) = \det(I - (g^{-1}e^{-\xi})^{\mathfrak{r}})$, hence

$$\kappa_g(D_{\mathfrak{k}}^g(N^g, \xi)) = \kappa_g(\det(I - (g^{-1}e^{-\xi})^{\mathfrak{r}})) D^{\Sigma}(N_{\Sigma}) | \Sigma_g. \quad (50)$$

With these preparations, we apply Theorem 3.2 to the integral 47, and obtain

$$I_g(m) = \sum_j' \frac{1}{d_j} \int_{\Sigma_j} \frac{Td(\Sigma_j) Ch^{\Sigma}(L_{\Sigma}^m)}{D^{\Sigma}(N_{\Sigma})} + O(m^{-\infty}), \quad (51)$$

the sum being over the connected components of Σ_g . Summing over all contributions, we get (15) up to an error term $O(m^{-\infty})$. As we remarked above, the right hand side of (15) is an arithmetic polynomial as a function of m . But if $f : \mathbb{N} \rightarrow \mathbb{Z}$ is an integer-valued function with $\lim_{m \rightarrow \infty} (f(m) - p(m)) = 0$ for some polynomial p , then $f(m) = p(m)$ for large m . This shows that the error term is zero for large m , and finishes the proof of the first part of Theorem 2.1.

5 Counting lattice points

To prove the second part of Theorem 2.1, i.e. that we can set $m = 1$, all we have to show is that $m \rightarrow N^{(m)}(m\mu)$ is an arithmetic polynomial.

Theorem 5.1 *Suppose that $J(M) \subset \mathfrak{g}_{reg}^*$. Then the function $m \mapsto N^{(m)}(m\mu)$ is an arithmetic polynomial for all $\mu \in \Lambda_+$.*

Before we prove this, we convert the computation of the multiplicities into a problem of counting lattice points. The next steps are based on work of Guillemin-Lerman-Sternberg [15] and Guillemin-Prato [16], except that we replace their use of the Atiyah-Bott Lefschetz Formula with the Localization Formula 3.1 applied to (37), since we do not want to assume isolated fixed points. Consider the action of the maximal torus $T \subset G$, with its moment map J^T equal to J followed by projection to $\mathfrak{t}^*$.

Proposition 5.1 *For all generic $\xi \in \mathfrak{t}$,*

$$\chi(e^\xi) = \sum_{\mathcal{F}} \int_{\mathcal{F}} \frac{Td(\mathcal{F}) e^{\omega + 2\pi i \langle J_{\mathcal{F}}, \xi \rangle}}{\det(I - e^{-\frac{i}{2\pi} F_{\mathfrak{t}}(N_{\mathcal{F}}, \xi)})}, \quad (52)$$

the sum being over the fixed points manifolds of the $T \subset G$ -action, $N_{\mathcal{F}}$ the corresponding normal bundles, and $J_{\mathcal{F}}$ the constant value of J on $\mathcal{F}$.

Proof. By “generic” we mean that the zero set of ξ_M is equal to the fixed point set of the T -action. To get (52), apply the Localization Formula, Theorem 3.1, to (37). The bundle $TM|_{\mathcal{F}}$ splits into the direct sum $T\mathcal{F} \oplus N_{\mathcal{F}}$. Since T does not act on $T\mathcal{F}$,

$$Td_{\mathfrak{t}}(M, \xi) = Td(\mathcal{F}) Td_{\mathfrak{t}}(N_{\mathcal{F}}, \xi).$$

The equivariant Euler class of $N_{\mathcal{F}}$ in Proposition 3.1 cancels the denominator of the equivariant Todd class, which immediately gives (52). $\square$

Although the left hand side of (52) is an analytic function of ξ , the individual summands on the right hand side have poles. Since they are not in L_{loc}^1 , they do not a priori define distributions on $\mathfrak{t}$. This problem can be fixed as follows [8]. By using the splitting principle (or simply a partition of unity on $\mathcal{F}$) if necessary, we can assume that $N_{\mathcal{F}}$ splits into a direct sum of invariant line bundles $N_{\mathcal{F}}^1, \dots, N_{\mathcal{F}}^r$. Let $\alpha_{\mathcal{F}}^j$ be the weight for the T -action on $N_{\mathcal{F}}^j$, that is, $e^\xi \in T$ acts by the character $\exp(2\pi i \langle \alpha_{\mathcal{F}}^j, \xi \rangle)$. Each $\alpha_{\mathcal{F}}^j$ determines an orthogonal hyperplane in $\mathfrak{t}$, let C be any fixed connected component in the complement of the union of all these hyperplanes. If we replace ξ by $\xi - i\eta$ in (52), with $\eta \in C$, the terms on the right hand side are analytic for all ξ . One can therefore regard (52) as an equality of distributions, with the summands on the right hand side defined as a distributional limit for $\eta \rightarrow 0$ in C .

Let us now first discuss the abelian case, i.e. assume that $G = T$ is a torus. Denote by $F^j(N_{\mathcal{F}})$ the components of the curvature. By expanding $\det(I - e^{-\frac{i}{2\pi}F_t(N_{\mathcal{F}}, \xi - i\eta)})^{-1}$ into its Taylor series w.r.t. $F^j(N_{\mathcal{F}})$, we can write it as a finite sum

$$\det(I - e^{-\frac{i}{2\pi}F_t(N_{\mathcal{F}}, \xi - i\eta)})^{-1} = \sum_{s \in \mathbb{N}^r} \frac{p_s(F^1(N_{\mathcal{F}}), \dots, F^r(N_{\mathcal{F}}))}{\prod_j (1 - e^{-2\pi i \langle \alpha_{\mathcal{F}}^j, \xi - i\eta \rangle})^{s_j}}, \quad (53)$$

where for all $s = (s_1, \dots, s_r)$, p_s is a polynomial. We now invoke the ‘‘polarization trick’’ used in ref. [15, 16]. For each $\alpha_{\mathcal{F}}^j$, write

$$\check{\alpha}_{\mathcal{F}}^j = \begin{cases} \alpha_{\mathcal{F}}^j & \text{if } \langle \alpha_{\mathcal{F}}^j, \eta \rangle > 0 \\ -\alpha_{\mathcal{F}}^j & \text{if } \langle \alpha_{\mathcal{F}}^j, \eta \rangle < 0 \end{cases} \quad (54)$$

for any, hence all, $\eta \in C$. Let $l_j^0 = 0$ if $\check{\alpha}_{\mathcal{F}}^j = \alpha_{\mathcal{F}}^j$, 1 otherwise. Then

$$\chi(e^{\xi - i\eta}) = \sum_{\mathcal{F}} \sum_{s \in \mathbb{N}^r} c_{\mathcal{F}, s} \frac{e^{2\pi i \langle J_{\mathcal{F}} - \sum l_j^0 s_j \check{\alpha}_{\mathcal{F}}^j, \xi - i\eta \rangle}}{\prod_j (1 - e^{-2\pi i \langle \check{\alpha}_{\mathcal{F}}^j, \xi - i\eta \rangle})^{s_j}} \quad (55)$$

with

$$c_{\mathcal{F}, s} = (-1)^{k_{\mathcal{F}, s}} \int_{\mathcal{F}} Td(\mathcal{F}) e^{\omega} p_s(F^1(N_{\mathcal{F}}), \dots, F^r(N_{\mathcal{F}})), \quad (56)$$

where $k_{\mathcal{F}, s} = \sum l_j^0 s_j$ is the number of sign changes.

For given $\mathcal{F}, s$, write $(a^1, \dots, a^N)$ for the list of $\check{\alpha}_{\mathcal{F}}^j$'s, appearing with respective multiplicities s_j . Since

$$(1 - e^{-2\pi i \langle a^j, \xi - i\eta \rangle})^{-1} = \sum_{l_j=0}^{\infty} e^{-2\pi i \langle l_j a^j, \xi - i\eta \rangle}, \quad (57)$$

we get

$$\chi(e^{\xi}) = \sum_{\mathcal{F}, s} c_{\mathcal{F}, s} \sum_{l \in \mathbb{Z}_+^N} e^{2\pi i \langle J_{\mathcal{F}} - \sum (l_j + l_j^0) a^j, \xi \rangle} \quad (58)$$

(the sum over $\mathbb{Z}_+^N := \{l \in \mathbb{Z}^N : l_j \geq 0\}$ is a well-defined periodic distribution). Comparing this to

$$\chi(e^{\xi}) = \sum_{\mu \in \Lambda} N(\mu) e^{2\pi i \langle \mu, \xi \rangle} \quad (59)$$

yields

$$N(\mu) = \sum_{\mathcal{F}, s} c_{\mathcal{F}, s} \mathfrak{P}_{\mathcal{F}, s}(J_{\mathcal{F}} - \mu - \sum l_j^0 a^j) \quad (60)$$

where the partition function $\mathfrak{P}_{\mathcal{F}, s}(\nu)$ is the number of solutions $k \in \mathbb{Z}^N$ of $\sum k_j a^j = \nu$, $k_j \geq 0$.

Starting from this expression, we will now show that $N^{(m)}(m\mu)$ is an arithmetic polynomial. We have to replace ω by $m\omega$, μ by $m\mu$ and J by mJ . Since $c_{\mathcal{F},s}^{(m)}$ is a polynomial in m , it is sufficient to show that the number of integer solutions of

$$m(J_{\mathcal{F}} - \mu) = \sum (l_j + l_j^0) a^j, \quad l_j \geq 0 \quad (61)$$

is an arithmetic polynomial as a function of m . Let us write $\nu = J_{\mathcal{F}} - \mu$, and consider $A = (a^1, \dots, a^N)$ as a $\mathbb{Z}$ -linear mapping $\mathbb{Z}^N \rightarrow \mathbb{Z}^p$, where $p = \dim(T)$. We are thus looking for integer solutions of

$$m\nu = Al, \quad l_j \geq l_j^0. \quad (62)$$

We will need the following

Theorem 5.2 (Ehrhart [11]) *Let L be a lattice, with underlying vector space $L_{\mathbb{R}} = L \otimes_{\mathbb{Z}} \mathbb{R}$, and $\Delta \subset L_{\mathbb{R}}$ a lattice polytope, i.e. a polytope whose vertices are all lattice points. Then, for all $r \in \mathbb{N}$, the counting function*

$$f(m) = \# \left(\frac{m}{r} \Delta \cap L \right) \quad (63)$$

is an arithmetic polynomial, with period r .

Let now $x_0 \in \mathbb{R}^N$ be any solution of $Ax = \nu$. The general solution of $Ax = m\nu$ is thus given by the affine plane $E_m = mx_0 + \ker(A)$. Let $r \in \mathbb{N}$ be the smallest number such that the vertices of the polytope $\Delta := E_r \cap \mathbb{R}_+^N$ are lattice points. If $l^0 = 0$, the set of solutions of (62) is the intersection $\frac{m}{r} \Delta \cap \mathbb{Z}^N$, so the number of solutions is an arithmetic polynomial by Ehrhart's Theorem. If $l^0 \neq 0$, let Δ_j be the face of Δ defined by $x_j = 0$, and let Δ' be the union of all Δ_j for which $l_j^0 = 1$. Then the solution set of (62) is

$$\frac{m}{r} \Delta \cap \mathbb{Z}^N - \frac{m}{r} \Delta' \cap \mathbb{Z}^N,$$

and this is again an arithmetic polynomial by Ehrhart's Theorem. This proves Theorem 5.1 in the abelian case.

Suppose now that G is nonabelian, but that $J(M)$ is contained in the set of regular elements, $\mathfrak{g}_{reg}^* = G \cdot \text{int}(\mathfrak{t}_+^*)$. We will show how this reduces to the abelian case. By the Symplectic Slice Theorem [18], $Y_+ = J^{-1}(\text{int}(\mathfrak{t}_+^*))$ is a symplectic (but not necessarily Kähler) submanifold of M , and is in fact a Hamiltonian T -space, with the restriction of J serving as a moment map. The above assumption implies that Y_+ is a *closed* submanifold, and $M = G \times_T Y_+$. The restriction $L_+ = L|_{Y_+}$ renders a quantizing bundle for Y_+ . Consider the expression

$$\chi'(e^\xi) := \int_{Y_+} Td_{\mathfrak{t}}(Y_+, \xi) Ch_{\mathfrak{t}}(L_+, \xi). \quad (64)$$

We claim that this is of the form

$$\chi'(e^\xi) = \sum_{\mu \in \Lambda} N'(\mu) e^{2\pi i \langle \mu, \xi \rangle}, \quad (65)$$

where $N'(\mu) \neq 0$ for only finitely many lattice points, and $N'(\mu) = 0$ unless $\mu \in J(Y_+) \cap \Lambda \subset \Lambda_+$. Indeed, one can check directly that $\chi(e^\xi)$ comes from a function on T , given near any point $g \in T$ by the formula (38), and then repeat the above analysis. (One can also pick a T -invariant almost Kähler structure on M , and then realize $\chi(e^\xi)$ as the equivariant index for some Dirac operator associated to the Clifford module $L \otimes \Lambda(T^{(0,1)}Y_+)^*$.)

Lemma 5.1 *For all $\mu \in \Lambda_+$, $N(\mu) = N'(\mu)$.*

Since we know that $N'^{(m)}(m\mu)$ is an arithmetic polynomial, this will finish the proof of Theorem 2.1.

Proof. Let us go back to the formula (52) for the character. Notice that the Weyl group $W = N_G(T)/T$ acts on M^T by permuting the connected components, and that M^T consists of its portion in Y_+ and the W -transforms thereof. Let $\mathcal{F} \subset Y_+$ be a connected component of M^T . The normal bundle $N_{\mathcal{F}}$ of $\mathcal{F}$ in M splits into its part in Y_+ , $N'_{\mathcal{F}} := N_{\mathcal{F}} \cap TY_+$, and the symplectic orthogonal of $TY_+|_{\mathcal{F}}$, which is canonically isomorphic to the trivial bundle $\mathfrak{g}/\mathfrak{t}$. The weights for the T -action on $\mathfrak{g}/\mathfrak{t}$ are of course simply the positive roots $\beta \in \mathfrak{t}^*$ of G . Therefore, by taking the trivial connection on $\mathfrak{g}/\mathfrak{t}$,

$$\det(I - e^{-\frac{i}{2\pi} F_t(N_{\mathcal{F}}, \xi)}) = \prod_{\beta > 0} (1 - e^{-2\pi i \langle \beta, \xi \rangle}) \det(I - e^{-\frac{i}{2\pi} F_t(N'_{\mathcal{F}}, \xi)}),$$

hence

$$\chi(e^\xi) = \sum_{w \in W} \frac{1}{\prod_{\beta > 0} (1 - e^{-2\pi i \langle \beta, w^{-1}(\xi) \rangle})} \sum_{\mathcal{F} \subset Y_+} \int_{\mathcal{F}} \frac{Td(\mathcal{F}) e^{\omega + 2\pi i \langle J_{\mathcal{F}}, w^{-1}(\xi) \rangle}}{\det(I - e^{-\frac{i}{2\pi} F_t(N'_{\mathcal{F}}, w^{-1}(\xi))})}.$$

To the sum

$$\sum_{\mathcal{F} \subset Y_+} \int_{\mathcal{F}} \frac{Td(\mathcal{F}) e^{\omega + 2\pi i \langle J_{\mathcal{F}}, \xi \rangle}}{\det(I - e^{-\frac{i}{2\pi} F_t(N'_{\mathcal{F}}, \xi)})},$$

we can apply the Localization Formula, this time in the opposite direction, and get that it is equal to the above expression $\chi'(e^\xi)$. This gives

$$\chi(e^\xi) = \sum_{w \in W} \frac{\chi'(e^{w^{-1}(\xi)})}{\prod_{\beta > 0} (1 - e^{-2\pi i \langle \beta, w^{-1}(\xi) \rangle})}.$$

But

$$\prod_{\beta > 0} (1 - e^{-2\pi i \langle \beta, w^{-1}(\xi) \rangle}) = \det(w) e^{-2\pi i \langle w(\delta) - \delta, \xi \rangle} \prod_{\beta > 0} (1 - e^{-2\pi i \langle \beta, \xi \rangle}),$$

where $\delta = \frac{1}{2} \sum_{\beta > 0} \beta$ is the magic weight. Weyl's Character Formula hence shows that

$$\chi(e^\xi) = \sum_{\mu \in \Lambda_+} N'(\mu) \sum_{w \in W} \det(w) \frac{e^{2\pi i \langle w(\delta + \mu) - \delta, \xi \rangle}}{\prod_{\beta > 0} (1 - e^{-2\pi i \langle \beta, \xi \rangle})} = \sum_{\mu \in \Lambda_+} N'(\mu) \chi_\mu(e^\xi),$$

where χ_μ is the character of the irreducible representation corresponding to μ . This proves $N(\mu) = N'(\mu)$. $\square$

Remarks.

1. If $J(M) \not\subset \mathfrak{g}_{reg}^*$, it is still possible to derive a formula for $N(\mu)$ similar to (60), following part II of Guillemin-Prato [16]. However, this formula involves an additional “shift”, so that (62) gets replaced by an equation of the form

$$Al = m\nu + \sigma, l_j \geq l_j^0.$$

In general, the number of integer solutions of such an equation is not an arithmetic polynomial for all $m \in \mathbb{N}$, even though this is true for large m .

2. On the other hand, Theorem 5.1 does not require that μ is a regular value of J . Even in the singular case, it is therefore sufficient to prove Multiplicity Formulas under the assumption $m \gg 0$.

Acknowledgements.

I would like to thank V. Guillemin, J. Kalkman, E. Lerman, R. Sjamaar and C. Woodward for useful comments and discussions. I am very much indebted to Victor Guillemin, whose recent results [14] on Multiplicity Formulas of Riemann-Roch type have been the basic motivation for this work. The stationary phase version (36) of the Jeffrey-Kirwan Localization Theorem was worked out jointly with Jaap Kalkman, who has also been a great help in explaining equivariant cohomology to me. This work was carried out when I was visiting scholar at the M.I.T., and I wish to thank the Mathematics Department for its hospitality.

Postscript: After completing this article, we learned about independent work of M. Vergne [24], who has made a different application of equivariant cohomology to the multiplicity problem.

References

- [1] M. F. Atiyah: Elliptic operators and compact groups. Lecture Notes in Mathematics **401**, Springer-Verlag (1974).

- [2] M. F. Atiyah, R. Bott: The moment map and equivariant cohomology. *Topology* **26**, 1-28 (1984).
- [3] M. F. Atiyah, G. Segal: The index of elliptic operators II. *Ann. Math.* **87**, 531-545 (1968).
- [4] M. F. Atiyah, I. Singer: The index of elliptic operators III. *Ann. Math.* **87**, 546-604 (1968).
- [5] N. Berline, M. Vergne: Classes caractéristiques équivariantes. Formules de localisation en cohomologie équivariante. *C. R. Acad. Sci. Paris* **295**, 539-541 (1982).
- [6] N. Berline, M. Vergne: The equivariant index and Kirillov's Character Formula. *Am. J. Math.* **107**, 1159-1190 (1985).
- [7] N. Berline, E. Getzler, M. Vergne: Heat kernels and Dirac operators, Springer-Verlag 1992.
- [8] J. J. Duistermaat: Equivariant cohomology and stationary phase. University Utrecht, Preprint 817, 1993.
- [9] J. J. Duistermaat, G. Heckman: On the variation in the cohomology of the symplectic form of the reduced phase space. *Inv. Math.* **69**, 259-268 (1982). Addendum, **72**, 153-158 (1983).
- [10] M. Duflo, M. Vergne: Cohomologie équivariante et descente. *Asterisque* **215** (1993).
- [11] E. Ehrhart: Polynômes arithmétiques et méthode des polyèdres en combinatoire. Birkhäuser Verlag, 1977.
- [12] C. Farsi: K- theoretical Index Theorems for orbifolds. *Quart. J. Math. Oxford* **43** (2), 183-200 (1992).
- [13] P. Griffiths, J. Harris: Principles of algebraic geometry. Wiley, 1978.
- [14] V. Guillemin: Reduction and Riemann-Roch. Preprint (1994).
- [15] V. Guillemin, E. Lerman, S. Sternberg: On the Kostant Multiplicity Formula. *J. Geom. Phys.* **5**, 721-750 (1988).
- [16] V. Guillemin, E. Prato: Heckman, Kostant and Steinberg Formulas for symplectic manifolds. *Adv. Math.* **82**, 160-179 (1990).
- [17] V. Guillemin, S. Sternberg: Geometric quantization and multiplicities of group representations. *Invent. Math.* **67**, 515-538 (1982).
- [18] V. Guillemin, S. Sternberg: Symplectic techniques in physics. Cambridge University Press 1984.

- [19] L. Hörmander: The analysis of linear partial differential operators I. Springer 1990.
- [20] L. Jeffrey, F. Kirwan: Localization for nonabelian group actions. Preprint (1993), alg-geom/9307001, to appear in Topology.
- [21] T. Kawasaki: The Riemann-Roch Theorem for complex V-manifolds. Osaka J. Math. **16**, 151-157 (1979).
- [22] I. Satake: The Gauss-Bonnet Theorem for V-manifolds. J. Math. Soc. Japan, **9**, 464-492 (1957).
- [23] R. Sjamaar: Holomorphic slices, symplectic reduction and multiplicities of representations (1993). To appear in Ann. Math.
- [24] M. Vergne: Quantification géométrique et multiplicités. Preprint (1994).